

Decade-Long Multi-Wavelength Study of Blazar OJ 287


Prelims: Current events of national and international importance | Science & Technology

Why in News?

Recently, the Department of Science & Technology (DST) announced key findings from a comprehensive decade-long observational study of the famous blazar OJ 287.

Key Observations

- The study tracked the multi-wavelength brightness, color variations, and spectrum of OJ 287.
- **Led by** - Scientists from the Aryabhata Research Institute of Observational Sciences (ARIES) in Nainital (in collaboration with the University of Turku, Finland, and international observatories).
- The research provides crucial confirmation regarding the dynamics of its **rare supermassive binary black hole (SMBH) system** and refines mass constraints on the primary black hole.
- **System Architecture** - Located roughly **4 billion light-years away** in the constellation Cancer, OJ 287 is an active galactic nucleus (AGN) belonging to the BL Lacertae (BL Lac) blazar class.
- It harbors a secondary supermassive black hole in a tight, precessing 12-year orbit around a primary black hole.
- **Mass Refinement of Central Black Hole** - By analyzing 8 optical spectra recorded during low-brightness states in 2017 using the **Steward Observatory (USA)**, researchers evaluated the spectral broadening of the O III **emission line**.
- This spectral analysis established that the central primary black hole's mass is **at least 3.89 billion solar masses** ($3.89 \times 10^9 M_{\odot}$).
- **Decade-Long Multi-Wavelength Flickering** - It recorded complex optical and X-ray variability.
- Helping untangle the complex interplay between thermal accretion disc emission, non-thermal jet radiation, and periodic impact flares produced when the smaller companion pierces the primary's accretion disc twice per orbit.



Significance

- **Testing General Relativity in Extreme Gravity Regimes**
 - OJ 287 acts as a cosmic laboratory to test Einstein's theory of **general relativity** in strong gravitational fields.
 - The companion's orbital precession rate is extreme ($\sim 39^\circ$ per orbit) due to energy loss from gravitational wave radiation, providing direct observational validation of general relativistic orbital mechanics.
- **Multi-Messenger Astronomy Target**
 - Supermassive binary black hole systems like OJ 287 are primary candidate sources for low-frequency **gravitational waves** detectable by Pulsar Timing Arrays (PTAs) and future space-based detectors like LISA.
 - Linking electromagnetic variations (light, X-rays) with gravitational wave predictions bridges key gaps in multi-messenger astrophysics.
- **Understanding Relativistic Jet Formation**
 - Insights into how dual black hole dynamics affect jet precession, particle acceleration, and energy output help explain how relativistic jets in active galaxies are powered and sustained.

Reference

[DST | Giant binary black hole system](#)



SHANKAR
IAS PARLIAMENT
Information is Empowering