

Glanders

Prelims - Current events of National and International importance | General issues on Environmental ecology, Bio-diversity.

Why in News?

Recently, the Department of Animal Husbandry & Dairying (DAHD), under the Ministry of Fisheries, Animal Husbandry & Dairying, issued a revised national action plan on glanders.

• Glanders is a highly contagious and often *fatal disease of equines*.

Equines are a **group of mammals** that belong to the family **Equidae.** Equines are known for their slender legs, hooves, and long faces with large, expressive eyes. Example <u>Horse, Donkey, Mules.</u>

- It can also affect other animals and humans.
- It was caused by the bacterium **Burkholderia mallei.**
- Clinical Signs It causes nodules and ulcerative lesions in the respiratory tract and lungs.
- A distinct cutaneous form, known as '<u>farcy'</u>, is also observed, marked by skin nodules and swollen lymph vessels.
- **Transmission** Ingestion is the primary route of infection, especially through shared feeding or watering sources.
- It is *transmitted to humans* through contact with tissues or body fluids of infected animals.
- The bacteria enter the body through cuts or abrasions in the skin and through mucosal surfaces such as the conjunctiva and nose.
- **Symptoms** Fever with chills and sweating, Muscle aches, Chest pain, Muscle tightness, Headache, Nasal discharge, Light sensitivity (sometimes with excessive tearing of the eyes).
- Global Occurrence It continues to appear sporadically in parts of Asia, Africa, the Middle East, and South American countries.
- Treatment Trimethoprim-sulfamethoxazole (TMP-SMX), A

sulphonamide combination.

- Piperacillin-tazobactam is being considered due to emerging resistance.
- Legal Frame work Recognized as a *notifiable disease* under the Prevention and Control of Infectious and Contagious Diseases in Animals (PCICDA) Act, 2009.
- It is a listed disease as described in the terrestrial animal health code of the World Organisation for Animal Health (WOAH).
- Recent Revised National Action Plan on Glanders
- Major Revisions
 - Reducing the infected zone from 5 km to 2 km,
 - Redefining the surveillance area as 2-10 km instead of 5-25 km, and
 - $_{\circ}$ Aligning the restrictions accordingly, which will now apply only up to 10 km.
- Enhanced Surveillance & Reporting.
- Strengthened Quarantine & Movement Control Rapid Response Mechanism.
- Capacity Building & Training.
- Public Awareness & Stakeholder Engagement.
- Research & Laboratory Support.

References

- 1. PIB| Revised Action Plan on Glanders
- 2. National centre for disease control | Glanders

