

# **Global Nutrition Report 2018**

#### What is the issue?

 $n\n$ 

\n

- The Global Nutrition Report 2018 was published recently.
- $\bullet$  With India facing a major malnutrition crisis, it calls for concerted actions to address the gaps and concerns.  $\mbox{\sc h}$

 $n\n$ 

### What are the key nutrition indicators?

 $n\n$ 

\n

- **Stunting**, or low height for age, is caused by long-term insufficient nutrient intake and frequent infections.
  - \n
- **Wasting**, or low weight for height, is a strong predictor of mortality among children under five.

\n

- $\bullet$  It is usually the result of acute significant food shortage and/or disease.
- **Overweight** is the condition of having more body fat than is healthy.  $\n$

 $n\n$ 

## What are the highlights of the report?

 $n\n$ 

\n

• Globally 150.8 million children under five years are stunted and 50.5 million are wasted.

\n

• Stunting - Of the three countries that are home to almost half (47.2%) of all stunted children, two are in Asia.  $\n$ 

• Of these, with 46.6 million (31%) children who are stunted, India tops the list of countries.

۱n

• It is followed by Nigeria (13.9 million) and Pakistan (10.7 million).

• **Wasting** - More than half of the world's children impacted by wasting (26.9 million) live in South Asia.

\n

- India accounted for 25.5 million children who are wasted, followed by Nigeria (3.4 million) and Indonesia (3.3 million).
- **Overweight** India also figures among the set of countries that has more than a million overweight children.

۱n

- $\bullet$  The other nations are China, Indonesia, Egypt, US, Brazil and Pakistan.  $\ensuremath{^{\backslash n}}$
- In four countries, more than a fifth of all children are overweight Ukraine, Albania, Libya and Montenegro.

\n

- $\bullet$  Of the around 38 million children globally overweight, some 5.4 million are in South Asia and 4.8 million are in East Asia. \n
- Prevalence of overweight children is the highest in upper-middle income countries and the lowest in low-income countries.

• In urban areas, there are 7.1% overweight children on average, whereas in rural areas 6.2% children are overweight.

۱'n

\n

- Overweight is slightly more common among boys (6.9%) than girls (6.1%).  $\$
- $\bullet$  As with obesity, among adults, women are more overweight than among men (39.2% and 38.5% respectively in 2016).  $\$
- $\bullet$  Conversely, diabetes is more common among men than women (9% and 7.9% respectively in 2014).

\n

- The health consequences of being overweight and obese contribute to an estimated four million deaths globally.
- **Malnutrition** Malnutrition is responsible for more ill-health than any other cause.

\n

• Of the 141 countries analysed, 88% of countries experience more than one form of malnutrition.

\n

- The problem of malnutrition remains severe across all regions but none of the countries are on course to meet all nine global nutrition targets.
- **Packaged foods** As per the report, only 21% of packaged foods in India are rated as being healthy.
- This is based on overall energy, salt, sugar and saturated fat on the negative side, and vegetable, fruit, protein, fibre and calcium as positive factors.

 $n\$ 

### What is the case with malnutrition in India?

 $n\n$ 

\n

• **Spatial variation** - An International Food Policy Research Institute (IFPRI) study found that stunting varied greatly from district to district (nearly 12% to 65%).

\n

- Notably, 239 of 604 districts have stunting levels above 40%.
- There are high and very high levels of stunting mainly in central and northern India (more than 30% and 40%, respectively).
- $\bullet$  Factors Factors associated with gender, education, economic status, health, hygiene, and other demographic factors explain the spatial variation. \n
- Factors such as women's low BMI accounted for 19% of the difference between the low versus high-burden districts.
- Other influential gender-related factors included maternal education (accounted for 12%), age at time of marriage (7%) and antenatal care (6%).
- Children's diets (9%), assets (7%), open defecation (7%) and household size (5%) were also influential.  $\n$

 $n\n$ 

### What should be done?

 $n\n$ 

\n

- The district- and region-wise sharp contrast shows the important role played by political commitment and administrative efficiency.
- It also reflects the role of literacy and women's empowerment in ensuring children's health.

۱'n

 $\bullet$  The figures thus call for immediate policy action to reduce inequalities and childhood stunting.  $\mbox{\sc h}$ 

 $n\n$ 

\n

• With district-wise data, state governments should address the determining factors for malnutrition.

\n

• Food and freedom go together, and the availability of one strongly influences access to the other.

\n

• So, social institutions can work to improve nutrition and children's welfare in free societies.

۱'n

- Governments should acknowledge the linkages and commit themselves to improved nutritional policies.
  - \n
- In states where the Anganwadi Services scheme does not work well, it must be subjected to a rigorous review, and targeted interventions for supplementary nutrition should be made.

 $n\n$ 

 $n\n$ 

**Source: Economic Times, The Hindu** 

\n

